

Claims

What is claimed is:

1. A method of identifying DNA sequences that control the expression of different collections of genes in a genome comprising, detecting selected DNA sequences adjacent to some genes excluding exons and introns.
2. A method of identifying DNA sequences that control the expression of different collections of genes comprising, detecting, by computer, one or more pairs of non-adjacent DNA sequences to which are bound to two RNA sequences.
3. A method of identifying DNA sequences that control the expression of different collections of genes in a genome comprising detecting changes in connectron behavior in the genome.
4. A method of modifying the expression of different gene collections in a genome, comprising detecting changes in connectron behavior as a result of an exogenous stimulus.
5. A method of detecting where and when new genes are being integrated into a host genome comprising detecting the connectrons in said host genome.
6. A method of detecting the expression effect of different gene collections in a given body comprising detecting the back and forth flow of connectrons between the chromosomes thereof.
7. A method of modifying a given body comprising modifying the connectron organization therein.

8. A method of detecting connectron control and target sequences in a given genome comprising:

determining the base composition of said genome,
determining one or more sites of control sequence organization, and/or
determining one or more sites of target application.

9. A method of determining the response of a cell in any tissue to changes in the cell's environment and/or genetic composition comprising providing a complete genomic DNA sequence for the organism and determining the effect of changes in connectrons due to application of a given exogenous stimulus to the genome.

10. In prokaryotes, archaea, single-celled eukaryotes and multi-celled eukaryotes, the tetradic relationship $T1=C1$ and $T2=C2$ where T1 and T2 are DNA sequences 20 or more bases in length, where the C1 sequence is adjacent to the C2 sequence, where the T1 and T2 sequences are on the same chromosome, and where the C1/C2 sequences are on the same chromosome as T1 and T2 or where the C1/C2 sequences are on a chromosome different from T1 and T2, wherein:

C1 sequence - any positive or negative strand DNA sequence of 20 bases or more, the C2 sequence must occur in the same chromosome as the C1 sequence,

C2 sequence - any positive or negative strand DNA sequence of 20 bases or more, the C1 sequence must occur in the same chromosome as the C2 sequence,

C1/C2 - any positive or negative strand DNA sequence of 40 or more bases such that the C1 sequence is adjacent to the C2 sequence,

T1 sequence - any positive or negative strand DNA sequence of 20 bases or more that is on the same chromosome as the T2 sequence, the T1 and T2 sequences must be between about 1kb and 105kb apart, and

T2 sequence - any positive or negative strand DNA sequence of 20 bases or more that is on the same chromosome as the T1 sequence, the T2 or T1 sequences must be between about 1kb and 105kb apart.

11. In prokaryotes, archea, single-celled eukaryotes and multi-celled eukaryotes, the connectron relationship that permits many different C1/C2 short loops to control the existence of a T1-T2 long loop and wherein said C1/C2 short lops can be on the same chromosome or on different chromosomes from the T1-T2 long loop, wherein:

C1 sequence - any positive or negative strand DNA sequence of 20 bases or more, the C2 sequence must occur in the same chromosome as the C1 sequence,

C2 sequence - any positive or negative strand DNA sequence of 20 bases or more, the C1 sequence must occur in the same chromosome as the C2 sequence,

C1/C2 - any positive or negative strand DNA sequence of 540 or more bases such that the C1 sequence is adjacent to the C2 sequence,

T1 sequence - any positive or negative strand DNA sequence of 20 bases or more that is on the same chromosome as the T2 sequence, the T1 and T2 sequences must be between about 1kb and 105kb apart, and

T2 sequence - any positive or negative strand DNA sequence of 20 bases or more that is on the same chromosome as the T1 sequence, the T2 or T1 sequences must be between about 1kb and 105kb apart.

12. In prokaryotes, archaea, single-celled eukaryotes and multi-celled eukaryotes, the connectron relationship that permits one C1/C2 short loop to control the existence of many T1-T2 long loops, the C1/C2 short loop can be on the same chromosome or on different chromosomes from the T1-T2 long loops, wherein:

C1 sequence - any positive or negative strand DNA sequence of 20 bases or more, the C2 sequence must occur in the same chromosome as the C1 sequence,

C2 sequence - any positive or negative strand DNA sequence of 20 bases or more, the C1 sequence must occur in the same chromosome as the C2 sequence,

C1/C2 - any positive or negative strand DNA sequence of 40 or more bases such that the C1 sequence is adjacent to the C2 sequence,

T1 sequence - any positive or negative strand DNA sequence of 20 bases or more that is on the same chromosome as the T2 sequence, the T1 and T2 sequences must be between about 1kb and 105kb apart, and

T2 sequence - any positive or negative strand DNA sequence of 20 bases or more that is on the same chromosome as the T1 sequence, the T2 or T1 sequences must be between about 1kb and 105kb apart.

13. The connectron relationships between prokaryotes and their plasmids wherein said connectrons implement a control mechanism between the two genomes that makes it possible from them to form a symbiotic relationship, and in the case of *D. radiodurans* the relationship is not symmetric, and the *D. radiodurans* genome sends C1/C2 short loops to the MP1 plasmid, wherein:

C2 sequence - any positive or negative strand DNA sequence of 20 bases or more, the C1 sequence must occur in the same chromosome as the C2 sequence,

C1/C2 - any positive or negative strand DNA sequence of 40 or more bases such that the C1 sequence is adjacent to the C2 sequence,

T1 sequence - any positive or negative strand DNA sequence of 20 bases or more that is on the same chromosome as the T2 sequence, the T1 and T2 sequences must be between about 1kb and 105kb apart,

T2 sequence - any positive or negative strand DNA sequence of 20 bases or more that is on the same chromosome as the T1 sequence, the T2 or T1 sequences must be between about 1kb and 105kb apart, and

3'UTR - untranslated 3' end of an mRNA is beyond the end of the last exon, a stop codon in the mRNA causes the ribosome to stop the translation of mRNA into protein.

16. In prokaryotes, archaea, single-celled eukaryotes and multi-celled eukaryotes, the connectron relationship that permits one C1/C2 short loop to control the existence of one or more T1-T2 long loops such that this C1/C2 short loop is itself subject to expression control by another T1-T2 long loop which surrounds it, wherein:

C1 sequence - any positive or negative strand DNA sequence of 20 bases or more, the C2 sequence must occur in the same chromosome as the C1 sequence,

C2 sequence - any positive or negative strand DNA sequence of 20 bases or more, the C1 sequence must occur in the same chromosome as the C2 sequence,

C1/C2 - any positive or negative strand DNA sequence of 40 or more bases such that the C1 sequence is adjacent to the C2 sequence,

T1 sequence - any positive or negative strand DNA sequence of 20 bases or more that is on the same chromosome as the T2 sequence, the T1 and T2 sequences must be between about 1kb and 105kb apart, and

T2 sequence - any positive or negative strand DNA sequence of 20 bases or more that is on the same chromosome as the T1 sequence, the T2 or T1 sequences must be between about 1kb and 105kb apart.

17. In prokaryotes, archaea, single-celled eukaryotes and multi-celled eukaryotes, the connectron relationship that permits one C1/C2 short loop to control the existence of the T1-T2 long loop that surrounds it, wherein:

C1 sequence - any positive or negative strand DNA sequence of 20 bases or more, the C2 sequence must occur in the same chromosome as the C1 sequence,

C2 sequence - any positive or negative strand DNA sequence of 20 bases or more, the C1 sequence must occur in the same chromosome as the C2 sequence,

C1/C2 - any positive or negative strand DNA sequence of 50 or more bases such that the C1 sequence is adjacent to the C2 sequence,

T1 sequence - any positive or negative strand DNA sequence of 20bases or more that is on the same chromosome as the T2 sequence, the T1 and T2 sequences must be between about 1kb and 105kb apart, and

T2 sequence - any positive or negative strand DNA sequence of 20 bases or more that is on the same chromosome as the T1 sequence, the T2 or T1 sequences must be between about 1kb and 105kb apart.

18. The connectron relationship that do not have any genes within the T1-T2 long loop, wherein:

T1 sequence is any positive or negative strand DNA sequence of 20 bases or more that is on the same chromosome as the T2 sequence, and

T2 sequence - any positive or negative strand DNA sequence of 20 bases or more that is on the same chromosome as the T1 sequence, and the T2 or T1 sequences must be between about 1kb and 105kb apart.

19. The geneless connectron relationship where one C1/C2 short loop controls the existence of many geneless T1-T2 long loops, wherein:

C1 sequence - any positive or negative strand DNA sequence of 20 bases or more, the C2 sequence must occur in the same chromosome as the C1 sequence,

C2 sequence - any positive or negative strand DNA sequence of 20 bases or more, the C1 sequence must occur in the same chromosome as the C2 sequence,

C1/C2 - any positive or negative strand DNA sequence of 40 or more bases such that the C1 sequence is adjacent to the C2 sequence,

T1 sequence - any positive or negative strand DNA sequence of 20 bases or more that is on the same chromosome as the T2 sequence, the T1 and T2 sequences must be between about 1kb and 105kb apart, and

T2 sequence - any positive or negative strand DNA sequence of 20 bases or more that is on the same chromosome as the T1 sequence, the T2 or T1 sequences must be between about 1kb and 105kb apart.

09036925 05-10-11